

# MerinoLink Limited Sire Evaluation

## Central Test Sire Evaluation *Within Flock Analysis*

### 2015 Drop Yearling Assessments

Conducted by



Engagement - Education  
Exploration - Exchange

under the auspices of

**The Australian Merino Sire Evaluation Association**



May 2016


# Moses & Son



## Wool • Merchandise • Livestock

Contact your local store to find out how we can meet your needs.

Temora:	02 6977 3100	Narrandera:	02 6959 1209
Wagga Wagga:	02 6922 7077	Young:	02 6382 5340
West Wyalong:	02 6972 4162	Ariah Park:	0400 965 532
Condobolin:	02 6895 2851	Adelaide:	0409 177 094





[www.mosesandson.com.au](http://www.mosesandson.com.au)



A Livestock Exchange like no other



## Informed Choices for your Sheep Marketing



Whether you're looking to buy or sell sheep, we can add value to your enterprise.

Call Simon Coddington on 02 6382 5340 or 0458 826 464  
Email: [simon@mosesandson.com.au](mailto:simon@mosesandson.com.au)

## MerinoLink Sire Evaluation Sponsors and Supporters – 2015 drop



### Acknowledgements

Dean Bourlet, Wynwood, Jugiong  
Mandy Bourlet, Wynwood, Jugiong  
Adele Offley, Moses & Son Young  
Simon Coddington, Moses & Son, Young  
Craig Wilson, Craig Wilson & Associates, Wagga

Jess Smith, Sally Martin Consulting, Young  
Mick Corkhill, Grassy Creek Merino Stud, Reids Flat  
Ben Patrick, Yarrawonga Merino Stud, Harden  
Lexi Cesnik, Sally Martin Consulting, Young

### Disclaimer

Australian Merino Sire Evaluation Association Incorporated (AMSEA) is funded by Australian Wool Innovation Limited (AWI) which gratefully acknowledges the funds provided by the Australian Government to support research, development and marketing of Australian wool. AMSEA sponsors, wool grower entry fees and site committee in-kind contributions also contribute to AMSEA funding. The print of this publication is funded by various advertisers and we thank them for their support. This publication should only be used as a general aid and is not a substitute for specific advice. To the extent permitted by laws AWI and AMSEA exclude all liability for loss or damage arising from the use of the information in this publication. © 2016 Australian Wool Innovation Limited and Australian Merino Sire Evaluation Association Incorporated. All rights reserved.

The Australian Merino Sire Evaluation Association has approved the format used in this report. Australian Flock Breeding Values reported here are based on analysis conducted by Sheep Genetics.

## MerinoLink Limited - Central Test Sire Evaluation

MerinoLink Limited run the South West Slopes sire evaluation site following the success of the South West Slopes Merino Breeders (2003, 2005 and 2008) and Bluechip Livestock (2011 x 2 and 2012) sire evaluations and young sire programs.

The MerinoLink Sire Evaluation site is an accredited Central Test Sire Evaluation (CSTE) site. It conforms to the requirement of the Australian merino Sire Evaluation Association (AMSEA).

We would like to thank and acknowledge the dedication of Dean and Mandy Bourlet for hosting the sire evaluation. Your enthusiasm and commitment to Merino breeding is appreciated and infectious.

The classing for the first visual assessments was conducted by Mr Mick Corkhill, Grassy Creek Merino Stud and Ben Patrick, Yarrawonga Merino Stud. We would like to fully acknowledge both Mick's and Ben's professional contribution to the visual assessments.

The 17 Merino sires being evaluated includes a historical sire, 1989 drop, as part of the AMSEA Historical Sire Program (funded by Australian Wool Innovation). The Historical Sire program

aims to evaluate sires that have been previously entered in sire evaluation sites between 15 and 20 years ago. The progeny will be fully evaluated alongside the current industry sires.

Whilst providing a very interesting comparison between leading sire of the 80's and 90's with those of today, the main purpose of the Historical Sire Program is to provide further validation and confidence in the system of linkage that is used by MERINOSELECT to directly compare animals across drops.

Results from the Historical Sire program will be published by AMSEA at the conclusion of the program in addition to individual sire performance results for historical sires being published in the MerinoLink Sire Evaluation Site Reports and the annual Merino Superior Sires.

We trust that everyone has achieved something out of this program and we look forward to providing leading genetic evaluation tools into the future.

**Sally Martin, MerinoLink CEO**

### Site Contacts

Name	Phone	Role
Dean Bourlet	0429 866 258	Host Property
Craig Wilson	0428 250 982	MerinoLink Board Director ( <i>Service Provider</i> )
Sally Martin	0400 782 477	Site Coordinator; Data Management; Reporting; MerinoLink Board Director ( <i>Service Provider</i> ); MerinoLink CEO
Richard Keniry	0427 878 541	MerinoLink Board Director ( <i>Commercial Breeder</i> )
Marty Moses	0417 691 308	MerinoLink Board Director ( <i>Service Provider</i> )
Carol Huggins	0429 934 616	MerinoLink Board Director ( <i>Ram Breeder</i> )
Steve Jarvis	0427 853 528	MerinoLink Board Director ( <i>Commercial Breeder</i> )
Robert Mortimer	02-6892 8259	MerinoLink Board Director ( <i>Ram Breeder</i> )
Mal Peake	0408 426 103	MerinoLink Board Director ( <i>Ram Breeder</i> )
Rick Baldwin	0429 833 837	MerinoLink Board Director ( <i>Ram Breeder</i> )
David Davidson	0429 847 345	MerinoLink Board Director ( <i>Commercial Breeder</i> )

### For further information on this report please contact:

Sally Martin, Sally Martin Consulting; 288 Maimuru Road, YOUNG NSW 2594;  
Mobile: 0400 782 477 Email: [sallymartin777@gmail.com](mailto:sallymartin777@gmail.com)

### Report authors:

Sally Martin<sup>1</sup> and Andrew Swan<sup>2</sup>

<sup>1</sup> Sally Martin Consulting, 288 Maimuru Road, YOUNG NSW 2594

<sup>2</sup> AGBU, UNE, Armidale, NSW 2351

**Date of publication:** May 2016

The information in this site evaluation report provides a comprehensive assessment of the 2015 drop 1<sup>st</sup> stage assessments of the sire's progeny performance, both measured and visually assessed traits. **The information reported is based on a within flock analysis of the sire progeny being evaluated.**

The 1<sup>st</sup> Assessment was carried out at 10 months of age with 10 months of wool growth. The 2<sup>nd</sup> Assessment is planned to be carried out at 18 to 22 months of age with 8 to 12 months wool growth.

## Contents

### Page

<b>Sire details</b>	4
<b>Management report</b>	5
<b>Assessment and management program</b>	6
<b>Visual trait assessment and site Breeding Objective</b>	7
<b>Results – 1<sup>st</sup> and 2<sup>nd</sup> Assessment</b>	
Summary	8
Figure 1a: Combined measured and visual trait performance (DPP)	8
Figure 1b: Combined measured and visual trait performance (MPP)	8
Figure 1c: Combined measured and visual trait performance (FPP)	8
Table 1: Index values and Classer's Grades	9
Figure 2: Fleece Weight and Fibre Diameter	10
Figure 3: Classer's Grade: Tops and Culls - Hogget	10
Figure 4: Fleece Weight and Body Weight	11
Detail	13
<b>Understanding the results – Measured trait performance</b>	13
Table 2: Major measured trait and Classer's Grade performance	14
Table 3: Other measured trait breeding values	15
<b>Understanding the results – Scored trait performance</b>	16
Table 4a: Wool quality	17
Table 4b: Wool quality & Pigmentation	18
Table 4c: Conformation	19
Table 4d: Breech	20
<b>Other assessment results</b>	
Table 5: Sire averages for measured traits	21
<b>Understanding the results – Information to assist the use of results</b>	
Index options	22
Accuracy of Flock Breeding Values (FBVs)	23
Link Sires	23
Calculation of combined information	23

## Sire and owner details

### Sire and owner details

Sire code	Breeders flock, Sire number Sire ID #, Breed <sup>†</sup>	Contact name, address Phone, Fax, Email
1* (Link)	Billandri Poll, 121391 600571-2012-121391 Poll Merino	Bill and Geoff Sandiland, Billandri, Kendenup WA 6323 P: (08) 9851 4030 F: (08) 9851 4264 E: <a href="mailto:csandilands@bordnet.com.au">csandilands@bordnet.com.au</a>
2	Bundilla, 120013 504081-2012-120013 Poll Merino	Rick Baldwin, Bundilla, 706 Tubbul Road, Young NSW 2594 P: (02) 6383 3802 F: (02) 6383 3837 E: <a href="mailto:bundillamerinos@bigpond.com">bundillamerinos@bigpond.com</a>
3*** Historic	Centre Plus Poll 000183 601250-1989-000183 Poll Merino	Robert Mortimer, Devondale, Tullamore NSW 2874 P: (02) 6892 8259 F: (02) 6892 8292 E: <a href="mailto:robert@centreplus.com.au">robert@centreplus.com.au</a>
4	Centre Plus Poll, 307564 601250-2013-307564 Poll Merino	Robert Mortimer, Devondale, Tullamore NSW 2874 P: (02) 6892 8259 F: (02) 6892 8292 E: <a href="mailto:robert@centreplus.com.au">robert@centreplus.com.au</a>
5**	Hazeldean, 003542 500383-2011-003542 Merino	Jim Litchfield, Hazeldean, Cooma NSW 2630 P: (02) 6453 5555 F: (02) 6453 2226 E: <a href="mailto:litchfield@hazeldean.com.au">litchfield@hazeldean.com.au</a>
6**	Kerin Poll, 130980 601413-2013-130980 Poll Merino	Nigel Kerin, Karuga Park, 1142 Bournewood Road, Yeoval NSW 2868 P: (02)6846 4070 E: <a href="mailto:kerinag@bigpond.com">kerinag@bigpond.com</a>
7**	Mumblebone, 130389 500063-2013-130389 Merino	Chad Taylor, Marrapana 456 Wuuluman Road, Wellington NSW 2820 P: (02) 6845m 3620 F: (02) 6845 3608 E: <a href="mailto:chad@mumblebone.com.au">chad@mumblebone.com.au</a>
8**	Mumblebone, 130850 500063-2013-130850 Merino	Chad Taylor, Marrapana 456 Wuuluman Road, Wellington NSW 2820 P: (02) 6845m 3620 F: (02) 6845 3608 E: <a href="mailto:chad@mumblebone.com.au">chad@mumblebone.com.au</a>
9	Pastora Poll, 130011 601090-2013-130011 Poll Merino	Tim Westblade, Pastora, Lochart NSW 2656 M: 0429 205 423 E: <a href="mailto:trwesty@bigpond.com">trwesty@bigpond.com</a>
10	Pastora Poll, 131634 601090-2013-131634 Poll Merino	Tim Westblade, Pastora, Lochart NSW 2656 M: 0429 205 423 E: <a href="mailto:trwesty@bigpond.com">trwesty@bigpond.com</a>
11	Pooginook Poll, 130083 601442-2013-130083 Poll Merino	John Sutherland, Paraway Pastoral Company, Jerilderie NSW 2716 P: (02) 6954 6145 F: (02) 6954 6168 E: <a href="mailto:pooginook@parawaypastoral.com">pooginook@parawaypastoral.com</a>
12	Rocklyn, 130022 501039-2013-130022 Merino	Ralph Diprose, Elon, Cowra Road, Grenfell NSW 2810 P: (02)6343 6331 F: (02) 6343 6331 E: <a href="mailto:rkdiprose@gmail.com">rkdiprose@gmail.com</a>
13**	Roseville Park, 140019 504166-2014-14001 Merino	Matthew Coddington, Glenwood, 39R Dilladerry Rd, Dubbo NSW 2830 P: (02) 6887 7286 F: (02) 6887 7103 E: <a href="mailto:rpmerinos@bigpond.com">rpmerinos@bigpond.com</a>
14	Centre Plus Poll (WA) 337919 609182-2013-337919 Poll Merino	Dave Vanenberghe, PO BOX 11 Scaddan WA 6447 P: (08) 9078 6049 M: 0427 786 049 E: <a href="mailto:riverlandwest@westnet.com.au">riverlandwest@westnet.com.au</a>
15	Wattle Dale 130115 503358-2013-130115 Merino	Dave Vanenberghe, PO BOX 11 Scaddan WA 6447 P: (08) 9078 6049 M: 0427 786 049 E: <a href="mailto:riverlandwest@westnet.com.au">riverlandwest@westnet.com.au</a>
16* (Link)	Willandra Poll, 120026 601610-2012-120026 Poll Merino	Ross Wells, Willandra, 477 North Coree Road, Jerilderie NSW 2711 P: (03) 5886 1223 F: (03) 5886 1605 E: <a href="mailto:rossirene@reachnet.com.au">rossirene@reachnet.com.au</a>
17	Woodpark Poll, 130431 601151-2013-130431 Poll Merino	Stephen and Carol Huggins, Eurolie, HAY NSW 2711 P: (02) 6993 4616 F: (02) 6993 4122 E: <a href="mailto:woodparkpoll@bigpond.com">woodparkpoll@bigpond.com</a>



## Graph and Table Key


- \* Link sire: Sire evaluated to provide links between years and sites so that the all site results can be combined into a single report, e.g., *Merino Superior Sires*.
- \*\* Common sires (in addition to Link Sires) between this CTSE site and other sites.
- # Sire ID provides a unique number for all sheep. A sire ID has 16 digits.
  - 2 for the breed of the flock, e.g., Merino (50), Poll Merino (60), Dohne (51), SAMM (48), Afrino (AF)
  - 4 for flock code, AASMB Registered flock code or unregistered code.
  - 4 for year of drop.
  - 6 for tag number used in the breeder's records.
- † Breed of flock in which the sire was born.
- ^ **Historical Sire** evaluated under AMSEA's R&D project to validate the system of linkage in MERINOSELECT that has operated over the past 15-20 years. These sires were generally widely used 15-20 years ago and were selected for the R&D program based on their high ASBV accuracies. They demonstrate the progress the industry has made over that period.

Example 16 digit code:	50	-	4967	-	2009	-	090012
	Breed		Flock		Year of drop		On-farm ID

**Advert – Grassy Creek**

**Advert – CW&S**

## 1. Location



“Wynwood”, 3035 Jugiong Road, Jugiong is located 30km from Harden on the Jugiong/Harden Road or 6km from Jugiong.

“Wynwood” is a commercial farming operation operated by Dean and Mandy Bourlet.

## 2. Selection and mating

- 900 medium framed ewes with free growing soft handling wools and low wrinkle score were selected and classed to be free from visual and conformation faults.
- The ewes were mated by Artificial Insemination to 17 sires.
- The ewes were randomly allocated to each sire.
- The insemination program was conducted on 12<sup>th</sup> and 13<sup>th</sup> February, 2015.
- The insemination program was conducted by Allstock, Dubbo.
- 50 ewes were allocated to each sire entered.

## 3. Pregnancy and lambing

- Pregnancy scanning took place on 13<sup>th</sup> May 2015.
- Ewes were managed as one contemporary group until 10 days before lambing.
- Adequate pasture and a supplementary feeding program ensured that nutritional requirements were met during all stages of pregnancy.
- Sire groups lambed down in separate paddocks.
- Lambs were tagged (visual and electronic) and weighed within two weeks of lambing and groups brought together and boxed into one contemporary group of ewes and lambs.

## 4. Weaning and seasonal conditions


- The lambs were marked and visually scored on 4<sup>th</sup> September 2015.
- The lambs were weaned onto improved and native pastures on 23<sup>rd</sup> October, 2015.

## 5. Visual Assessments

- The 1<sup>st</sup> stage visual assessment was carried out by Mick Corkhill and Ben Patrick.

## 6. Rainfall – Young (closest weather station)

	2012	2013	2014	2015	2016	Median
Jan	30.2	7.0	16.0	55.6	92.2	26.0
Feb	168.6	42.6	35.0	30.6	21.6	32.6
Mar	162.6	60.6	132.8	5.2	27.8	30.2
Apr	15.2	12.2	50.0	72.8	21.6	22.2
May	37.0	28.0	37.6	39.6	111.2	28.0
Jun	56.6	108.8	67.4	37.0		52.4
Jul	62.4	58.2	41.4	84.2		55.8
Aug	45.2	36.8	26.0	67.8		48.4
Sep	35.8	27.8	26.2	11.0		40.6
Oct	14.0	20.0	22.6	7.6		44.6
Nov	41.6	26.8	16.2	145.2		50.6
Dec	43.0	16.0	55.0	35.2		40.2
Totals	712.2	444.8	526.2	526.2		613



\*Source: BOM - Median 1995-2016.



## Assessment and management program

Activity	Date/s	Age (months)	Wool (months)
Selection of ewes & allocation of ewes for mating	07.01.2015		
Artificial Insemination	11.02.2015		
	12.02.2015		
Pregnancy scanning	13.05.2015		
Separated into sire lambing groups	30.07.2015		
Lambing: start – finish	08 to 15.07.2015		
Lambing mobs boxed to 1 management group	01.08.2015	14-21 days	
Tagging/pigment scores (age in days)	01.08.2015	14-21 days	
Marked and scored for breech traits	04.09.2015	55 days	
Weaning (age in days)	23.10.2015	104 days	
Pre assessment (even-up) shearing	NA		
Crutching			
• Post Weaning (PW)	02.02.2015	6.5	6.5
Fat and eye muscle scanning			
• Yearling			
Fleece sampling assessment			
• Yearling (Y)	26.04.2016	10	10
• Hogget (H)			
Staple length assessment			
• Yearling (Y)	26.04.2016	10	10
Classer's Grade assessment			
• Yearling (Y)	26.04.2016	10	10
• Hogget (H)			
Pre shearing scoring assessment			
• Yearling (Y)	26.04.2016	10	10
• Hogget (H)			
Assessment shearing			
• Yearling (Y)	27.04.2016	10	10
• Hogget (H)			
Post shearing scoring assessment			
• Yearling (Y)	28.04.2016	10	0
• Hogget (H)			
Body weigh assessment			
• Weaning (W)	23.10.2015	4	
• Post Weaning (PW)	21.01.2016	7	
• Yearling (Y)	26.04.2016	10	
• Hogget (H)			
Worm egg count sampling	Little challenge to date; still to be measured.		
• Yearling (Y)			
Sire's Progeny Group Evenness assessment			
Vaccination	Marking, weaning, post shearings		
Drench	As required based on worm egg counts		
Supplementary feeding: start – finish			
Field day or public display of sheep	21.01.2016 (pre 1 <sup>st</sup> stage shearing)		

## Visual Trait Assessment and site Breeding Objective

### Visual trait assessment

#### 1<sup>st</sup> Stage Assessment (Yearling)

**Classer's Grade:** Mick Corkhill

**Trait Scores:** Mick Corkhill, Ben Patrick and Adele Offley

**Breech Scores:** Sally Martin

#### **Site Breeding Objective used to assess the Classer's Grades – 1<sup>st</sup> Stage Assessment**

The Breeding Objective used to select the Classer's Tops (11%), Flock (40%) and Cull (29%) was based on a visual assessment where the animal performed well for growth (meet minimum body weight suitable for joining), were structurally sound with good wool quality traits including long soft handling wool and fleece weight. *(No reference was made to measured performance at the time of classing and was based on the visual presentation of all traits).*

#### **Within Site Analysis**

This report provides information within site on the performance of the progeny of the sires being evaluated. The ASBVs have not been taken into consideration in the within site analysis, however will be used in the across site (MSS) analysis. The information presented is a reflection of one sires performance, not the bloodline.

Publication of results in both Merino Superior Sires (MSS) and MerinoSelect will be presented as across flock Australian Sheep Breeding Values (ASBV's) and will included additional data collected on farm, at other sire evaluation sites and the Information Nucleus Flock sites (Resource Flock).



*Adele Offley, Moses & Son*



*Foundation ewes - Sire Allocation (pre AI)*

Advancing technology...  
maximising productivity.



THE ALLFLEX RS420 STICK READER.

**ACCURATE DNA  
TESTING IS HERE!**



TSU APPLICATOR



TSU INDIVIDUAL UNIT

**NEW BLUETOOTH  
DOCKET PRINTER  
- SIMPLE TO OPERATE!**



**RapID  
TAG**



Over the past two years at Wanganella and Poll Boonoke Merino Studs, we have used over 10,000 RapIDTags. We have found them easy to put in lambs with excellent retention. Within our stud operations, we have had 100% read performance.

**Angus Munro**  
Stud Manager,  
Australian Food and Agriculture



Identify the Difference

Contact our friendly Customer Service Team  
Ph. 1300 138 247 / [www.allflex.com.au](http://www.allflex.com.au)

## **YARRAWONGA**

**MERINO & POLL MERINO**



*Semen Sire Yarra 961 (PP)*

On Property ram sale  
Monday 12th September 2016  
Cunningham Plains  
Harden NSW

**Steve and Liz Phillips**  
Phone: 02 6386 3128  
Mobile: 0427 863 128

**Ben Patrick**  
Mobile: 0428 984 699

E: [yarrowongamerino@bigpond.com](mailto:yarrowongamerino@bigpond.com)  
W: [www.yarrowongamerino.com.au](http://www.yarrowongamerino.com.au)



## Figure 1a, 1b and 1c. Combined measured traits and visual trait performance

### Summary graph: visual and measured performance

Each sire meeting the AMSEA index accuracy threshold assessed at 1<sup>st</sup> Assessment is located on Figure 1a, 1b and 1c and describes the performance for combined measured traits and combined visual assessment.

A different graph is provided for each of the three production indexes reported. In each graph the combined measured traits are based on the AMSEA index and the visual trait performance is a combination of Classer's Grade performance (Tops and Culls). More information is found in "Calculation of combined performance" (page 25).

Sires that are above average performers for combined measured traits and Classer's Grade are located in the top right hand quarter.

Sire code	Breeders flock, Sire number	Sheep Genetics ID	Sire of Sire
1*	Billandri Poll, 121391	600571-2012-121391	600571-2010-100011
2	Bundilla, 120013	504081-2012-120013	504081-2010-100027
3	Centre Plus Poll, 307564	601250-2013-307564	601250-2009-907538
4^	Centre Plus Poll, 9.183	601250-1989-000183	601250-1985-500279
5	Centre Plus WA Poll, 337919	609182-2013-337919	601250-2010-007257
6	Hazeldean, 003542	500383-2011-003542	601050-2002-020603
7	Kerin Poll, 130980	601413-2013-130980	601244-2007-070304
8	Mumblebone, 130389	500063-2013-130389	601365-2009-090399
9	Mumblebone, 130850	500063-2013-130850	500063-2010-100186
10	Pastora Poll, 130011	601090-2013-130011	
11	Pastora Poll, 131634	601090-2013-131634	601090-2011-113416
12	Pooginook Poll, 130083	601442-2013-130083	600815-2010-101508
13	Rocklyn, 130022	501039-2013-130022	501039-2011-110155
14	Roseville Park, 140019	504166-2014-140019	601050-2009-090853
15	Wattle Dale, 130115	503358-2013-130115	601250-2009-907538
16*	Willandra Poll, 120026	601610-2012-120026	600610-2010-100033 "The GP"
17	Woodpark Poll, 130431	601151-2013-130431	601151-2011-110178

\* Link sire: Sire evaluated to provide links between years and sites so that the all site results can be combined into a single report, e.g., *Merino Superior Sires*.

^ **Historical Sire** evaluated under AMSEA's R&D project to validate the system of linkage in MERINOSELECT that has operated over the past 15-20 years. These sires were generally widely used 15-20 years ago and were selected for the R&D program based on their high ASBV accuracies. They demonstrate the progress the industry has made over that period.

**Independent advice and services  
YOU CAN DEPEND ON**



- Electronic identification systems
- Sheep production
- Breeding program design
- Benchmarking
- Data Management and Analysis
- Provider of RAMPOWER Indexes
- Pedigree Match Maker Analysis

Working with commercial breeders, ram breeders and service providers to make a difference for the sheep industry!



**m: 0400 782 477**

**e: sallymartin777@gmail.com**

**w: www.sallymartinconsulting.com.au**

**Figure 1a. Combined visual and measured performance**




Figure 1a is based on an **AMSEA Dual Purpose Plus (DP+) index** – (Based on a meat focused production system where surplus progeny are sold as lambs and a portion of ewes are joined to terminal sires).

**Figure 1b. Combined visual and measured performance**




Figure 1b is based on an **AMSEA Merino Production Plus (MP+) index** – (Based on a balanced wool and meat production system where surplus progeny are sold as hoggets).

**Figure 1c. Combined visual and measured performance**




Figure 1c is based on an **AMSEA Fibre Production Plus (FP+) index** (Based on a wool focussed production system where wethers are retained, operating in an environment where worms cause economic losses).

**Table 1. AMSEA Index values and Classer's Grade**

The highest performing 3 sires for each trait (i.e. trait leaders) are highlighted by shading. Each sire is listed for Classer's Grade and the same three indexes at all site evaluations. The index values reported are based on Flock Breeding Values (within flock) measured trait performance with varying emphasis on fleece weight, fibre diameter, body weight, staple strength and worm egg count. (See 'Index Options' on page 24 for more information on the indexes presented in the table below.) **AMSEA Indexes are the same as MERINOSELECT Indexes apart from NLW (Number of Lambs Weaned) which is given a zero FBV value in AMSEA calculations.**

<b>Dual Purpose Plus (DP+)</b>	Based on a meat focused production system where surplus progeny are sold as lambs and a portion of ewes are joined to terminal sires.
<b>Merino Production Plus (MP+)</b>	Based on a balanced wool and meat production system where surplus progeny are sold as hoggets.
<b>Fibre Production Plus (FP+)</b>	Based on a wool focussed production system where wethers are retained, operating in an environment where worms cause economic loss

Ram code	Breeders flock, Ram number	No of Progeny	AMSEA Indexes values			Classer's Grade	
			Fibre Production Plus	Merino Production Plus	Dual Purpose Plus	Tops % (dev)	Culls % (dev)
						Y <sup>2</sup>	Y
1*	Billandri Poll, 121391	36	95	95	95	3	-1
2	Bundilla, 120013	33	102	102	103	7	-16
3	Centre Plus Poll, 307564	31	117	130	134	2	-12
4*	Centre Plus Poll, 9.183	27	96	98	97	-7	25
5	Centre Plus WA Poll, 337919	29	112	119	122	6	-4
6	Hazeldean, 003542	30	113	113	95	2	-5
7	Kerin Poll, 130980	45	109	119	115	0	-9
8	Mumblebone, 130389	40	77	62	67	-8	14
9	Mumblebone, 130850	38	99	99	111	0	-5
10	Pastora Poll, 130011	38	98	93	87	-1	7
11	Pastora Poll, 131634	15	99	97	102	-11	11
12	Pooginook Poll, 130083	34	104	109	106	-8	1
13	Rocklyn, 130022	29	98	95	91	6	2
14	Roseville Park, 140019	31	99	100	99	8	-6
15	Wattle Dale, 130115	27	98	93	96	-7	1
16*	Willandra Poll, 120026	35	93	91	90	0	-12
17	Woodpark Poll, 130431	29	91	85	91	6	8
Average performance		32	100	100	100	11	29




- \* Link sire: Sire evaluated to provide links between site evaluations and sites so that the all site results can be combined into a single report, e.g., *Merino Superior Sires*.
- 1 Classer's Grade is expressed as the percentage deviation of average Tops% and Culls%
- 2 Y = Yearling (300 to 400 days). H = Hogget (400 to 540 days)
- ^ **Historical Sire** evaluated under AMSEA's R&D project to validate the system of linkage in MERINOSELECT that has operated over the past 15-20 years. These sires were generally widely used 15-20 years ago and were selected for the R&D program based on their high ASBV accuracies. They demonstrate the progress the industry has made over that period.




*2015 Drop – Lamb Marking (Early Sept 2015)*




*2015 Drop – Weaning (Late October 2015)*



**Figure 2 Fleece weight by fibre diameter FBVs (Hogget)** – describes performance for fleece weight on the side axis and fibre diameter on the bottom axis. Sires that are above average for fleece weight and below average fibre diameter are located in the top left hand quarter.



**Figure 3 Classer's Tops by Cull Grade** – describes performance for Classer's Tops Grade on the side axis and Cull Grade on the bottom axis. Sires that have above average Tops and below average Culls are in the top left hand quarter. Classer's Tops (23%), Flock (47%) and Cull (30%) is based a visual assessment where the progeny performed well for growth, structurally sound with good wool quality traits including long soft handling wool and fleece weight.



**Figure 4. Fleece weight by body weight (FBV's)** – describes performance for fleece weight on the side axis and body weight on the bottom axis. Sires that are above average for fleece weight and above average for body weight are located in the top right hand quarter.

## Understanding the results – measured trait performance

Measured trait performance and Classer's Grade – Tables 2 and 3 – pages 16 and 17.

<b>Sire code:</b>	Allows a sire to be located on the summary graphs and some tables.	
<b>Sire name:</b>	Identity of the breeder's flock and the sire's number or name.	
<b>Number of progeny:</b>	The number of progeny a sire had at the most recent measured analysis.	
<b>Horn/Poll:</b>	The Poll test has been developed by the Sheep CRC using measurements and DNA tests on animals in the Information Nucleus Flocks. The test is based on two genetic markers that are very close to the Poll gene. PP = Polled; PH = Half Poll; HH = Horned; blank = test failed	
<b>Flock Breeding Values:</b>	Flock Breeding Values (FBVs) are Estimated Breeding Values (EBVs) calculated by Sheep Genetics for the sire's evaluated in this report. Only data from this site evaluation is used in the calculation of these FBVs. FBVs describe the relative breeding value (genetic performance) of the sires (in this case based on the performance of their progeny). A sire's progeny will express half of their sire's FBV. FBVs do not necessarily reflect the sires observed performance, which is a combination of both genetic and environmental influences. FBVs are an estimate of the genetic component of the sheep's performance.	
<b>Traits:</b> Abbreviation, trait and the (units reported)	GFW: Greasy fleece weight (percentage). CFW: Clean fleece weight (percentage). FD: Average fibre diameter (micron). WT: Body weight (kilograms). FDCV: Fibre diameter coefficient of variation (percentage). SL: Staple length (mm) at the mid-side.	SS: Staple strength (N/ktex) at the mid-side. EMD: Eye muscle depth (mm) at the 'C' site. FAT: Fat depth (mm) at the 'C' site. CURV: Fibre curvature (degrees) WEC: Worm egg count (% deviation in worm burden of sire's progeny)
<b>Age at assessment:</b>	W = Weaning - 42 to 120 days (6 weeks to 4 months of age) E = Early Post Weaning - 120 to 210 days (4 to 7 months of age) P = Post Weaning - 210 to 300 days (7 to 10 months of age) Y = Yearling - 300 to 400 days (10 to 13 months of age) H = Hogget - 400 to 540 days (13 to 18 months of age) A = Adult - 540 days or older (18 months and older)	
<b>Classer's Grade:</b>	A Classer grades all progeny as either, Tops, Flocks or Culls based on their visual assessment of all traits relative to the site's Breeding Objective. The percentage deviation from the average of Tops and Culls is presented in this report. Average percentage of Tops and Culls for the entire drop is included in Table 1.	

Table 2. Major measured traits and Classer's Grades

Ram code      Breeders flock, Ram number		No.  of prog.	Flock Breeding Values (deviations)						Classer's Grade <sup>1</sup>	
			GFW %	CFW %	FD um	WT kg			Tops % (dev)	Culls % (dev)
						W	P	Y		
			Y <sup>2</sup>	Y	Y	W	P	Y	Y	Y
1*	Billandri Poll, 121391	36	1.0	1.0	-0.4	0.4	0.0	-0.3	3	-1
2	Bundilla, 120013	33	0.0	0.0	0.4	0.2	0.4	0.5	7	-16
3	Centre Plus Poll, 307564	31	12.0	12.0	-0.2	2.9	4.9	6.7	2	-12
4^	Centre Plus Poll, 9.183	27	3.0	2.0	0.4	-0.4	-0.8	-0.5	-7	25
5	Centre Plus WA Poll, 337919	29	6.0	5.0	-0.5	1.3	3.0	4.0	6	-4
6	Hazeldean, 003542	30	8.0	9.0	-1.0	-1.1	-2.4	-3.1	2	-5
7	Kerin Poll, 130980	45	13.0	16.0	0.9	1.0	2.0	2.5	0	-9
8	Mumblebone, 130389	40	-14.0	-16.0	0.6	-3.2	-4.3	-5.6	-8	14
9	Mumblebone, 130850	38	-5.0	-6.0	0.5	1.0	2.5	3.1	0	-5
10	Pastora Poll, 130011	38	-3.0	-3.0	-0.5	-0.7	-1.9	-3.0	-1	7
11	Pastora Poll, 131634	15	-6.0	-6.0	-0.6	0.2	0.7	0.9	-11	11
12	Pooginook Poll, 130083	34	7.0	9.0	0.9	0.7	0.9	1.1	-8	1
13	Rocklyn, 130022	29	-1.0	0.0	0.4	-0.9	-1.5	-2.3	6	2
14	Roseville Park, 140019	31	0.0	0.0	0.2	0.5	0.5	0.0	8	-6
15	Wattle Dale, 130115	27	-8.0	-8.0	-0.8	-0.6	-1.2	-0.9	-7	1
16*	Willandra Poll, 120026	35	-6.0	-4.0	0.3	-0.2	-1.1	-1.8	0	-12
17	Woodpark Poll, 130431	29	-8.0	-11.0	-0.7	-1.3	-1.3	-1.3	6	8
Average performance		32	3.0	2.0	16.4	24.9	33.2	37.0	11	29
			kg	kg	um	kg	kg	kg	%	%

\* Link sire: Sire evaluated to provide links between site evaluations and sites so that the all site results can be combined into a single report, e.g., *Merino Superior Sires*.

<sup>1</sup> Classer's Grade is expressed as the percentage deviation of average Tops% and Culls%

<sup>2</sup> Y = Yearling (300 to 400 days). H = Hogget (400 to 540 days)

<sup>^</sup> **Historical Sire** evaluated under AMSEA's R&D project to validate the system of linkage in MERINOSELECT that has operated over the past 15-20 years. These sires were generally widely used 15-20 years ago and were selected for the R&D program based on their high ASBV accuracies. They demonstrate the progress the industry has made over that period.

Information on how to use the results in the table above can be found on page 15.

Table 3. Other measured traits

Ram code	Breeders flock, Ram number	No. of prog.	Flock Breeding Values (deviations)						
			FDCV %	SL mm	SS N/ktex	Curv deg/mm	Fat mm	EMD mm	WEC%
			Y <sup>2</sup>	Y	Y	Y	Y	Y	Y
1*	Billandri Poll, 121391	36	0.6	-1.5	-4.7	4.6	Trait not measured at time of publication	Trait not measured at time of publication	Trait not measured at time of publication
2	Bundilla, 120013	33	-0.9	2.6	3.7	-0.6			
3	Centre Plus Poll, 307564	31	-0.6	3.1	-0.5	-0.1			
4^	Centre Plus Poll, 9.183	27	0.2	-6.7	0.2	4.3			
5	Centre Plus WA Poll, 337919	29	1.0	1.3	-1.3	-0.3			
6	Hazeldean, 003542	30	1.2	-4.4	-1.4	-0.2			
7	Kerin Poll, 130980	45	-0.1	9.2	3.2	-7.3			
8	Mumblebone, 130389	40	-1.4	5.5	0.2	-3.9			
9	Mumblebone, 130850	38	-1.9	8.1	3.1	-2.5			
10	Pastora Poll, 130011	38	0.7	-4.3	-0.7	3.5			
11	Pastora Poll, 131634	15	0.5	-2.3	-2.8	4.2			
12	Pooginook Poll, 130083	34	-0.3	10.7	4.1	-9.3			
13	Rocklyn, 130022	29	-0.5	2.6	2.7	-3.9			
14	Roseville Park, 140019	31	0.0	-1.4	0.6	-1.9			
15	Wattle Dale, 130115	27	0.3	-7.6	-2.4	6.8			
16*	Willandra Poll, 120026	35	0.4	-7.4	1.0	0.9			
17	Woodpark Poll, 130431	29	0.7	-7.6	-4.8	5.8			
Average performance		32	18.8 %	78.9 mm	44.7 N/ktex	90.5 deg/mm	mm	mm	%

<sup>2</sup> W = Weaning (42 to 120 days); P = Post Weaning (120 to 300 days); Y = Yearling (300 to 400 days); H = Hogget (400 to 540 days); A = Adult (540 days and older).

<sup>^</sup> **Historical Sire** evaluated under AMSEA's R&D project to validate the system of linkage in MERINOSELECT that has operated over the past 15-20 years. These sires were generally widely used 15-20 years ago and were selected for the R&D program based on their high ASBV accuracies. They demonstrate the progress the industry has made over that period.

Information on how to use the results in the table above can be found on page 15.

## Understanding the results – scored performance traits

**Scored trait performance** – Tables 4a to 4d – pages 19 to 22. The following description of trait scores is a summary of the detailed word and diagrammatical description of these scores in the Visual Sheep Scores booklet (free on application to AWI or downloadable at [www.merinosuperiorsire.com.au](http://www.merinosuperiorsire.com.au)). A deviation from the average trait score for all progeny is reported as well as the percentage of the sire's progeny recorded for each trait.

■ Fleece rot:	The severity of fleece rot from <b>1</b> (no fleece rot), <b>2 and 3</b> (bands of bacterial staining but no crusting), and <b>4 and 5</b> (bands of crusty fleece rot).
■ Wool colour:	Greasy wool colour scored from <b>1</b> (whitest) to <b>5</b> (yellow).
■ Wool character:	Definition and variation of crimp between and along the staple scored from <b>1</b> (well defined and regular) to <b>5</b> (undefined and large variation).
■ Dust penetration:	Degree of dust penetration from <b>1</b> (only tip <5%) to <b>5</b> (80 to 100% of staple).
■ Staple weathering:	The deterioration due to light and water from <b>1</b> (least, <5% of staple) to <b>5</b> (most, 30 to 50%) reflect the depth and degree of deterioration.
■ Staple structure:	The size and diameter of each staple from <b>1</b> (<5mm) to <b>5</b> (30 to 50 mm)
■ Face cover:	Wool cover on the face scored from <b>1</b> (open face) to <b>5</b> (fully covered face).
■ Feet/Legs:	Conformation of feet and legs scored from <b>1</b> (very good) to <b>5</b> (very poor).
■ Body wrinkle:	The degree of body wrinkle from <b>1</b> (no wrinkle) to <b>5</b> (extensive wrinkle).
■ Jaw:	Under- or over-shot lower jaw (and teeth) relative to the top jaw. Three scores: <b>1</b> (very well aligned), <b>3</b> (marginally under or over) and <b>5</b> (heavily under or over).
■ Back/Shoulder:	Conformation of the back and shoulder from <b>1</b> (very good) to <b>5</b> (very poor).
■ Fibre pigmentation:	The percentage of dark fibres on any part of the sheep from <b>1</b> (0 pigmented fibres at any site) to <b>5</b> (76 to 100% pigmented fibres at one or more sites). This trait does not include random spot or recessive black.
■ Non-fibre pigmentation:	The percentage of pigmentation on the areas not shorn from <b>1</b> (0 pigmentation at any site) to <b>5</b> (76 to 100% pigmented area on one or more bare skin sites, <b>and/or</b> 76 to 100% of the total hoof area).
■ Recessive black: (black)	Recessive black (black) is identified by relatively symmetrical markings on both sides of the face. There are two scores <b>1</b> (no recessive markings) and <b>5</b> (recessive markings). This trait does not include random spot or fibre pigmentation.
■ Random spot: (spot)	Random spot (spot) is identified by rounded wool or hair spot/s, not symmetrical. There are two scores <b>1</b> (no spot/s) and <b>5</b> (spot/s). If both sides of the face or body are spotted the sheep should be scored as a recessive black.
■ Breech cover	Size of natural bare area around the breech from <b>1</b> (large) to <b>5</b> (no bare).
■ Crutch cover	Size of natural bare area in the pubic and groin region from <b>1</b> (large) to <b>5</b> (no bare).
■ Breech wrinkle	Degree of wrinkle at the tail set and hind legs from <b>1</b> (nil) to <b>5</b> (extensive).
■ Dag	Degree of dag adhering to the breech and legs from <b>1</b> (nil) to <b>5</b> (extensive).
■ Urine	Degree of urine stained wool in the breech area, including the hind legs from <b>1</b> (nil) to <b>5</b> (extensive).



**Table 4a. Visual trait assessments – Wool quality**

Traits are reported as a deviation (Dev) from the average trait score for all progeny. The percentage of a sire's progeny assessed for each score is also reported. For the majority of breeder's objectives a negative deviation would be considered favourable and the larger the deviation the better.

Breeders flock, Sire number	Wool Quality																							
	Fleece Rot						Wool Colour						Wool Character						Dust Penetration					
	Dev	1	2	3	4	5	Dev	1	2	3	4	5	Dev	1	2	3	4	5	Dev	1	2	3	4	5
Billandri Poll, 121391	-0.1	97	3	0	0	0	0.0	36	61	3	0	0	0.0	42	50	8	0	0	0.0	0	58	42	0	0
Bundilla, 120013	-0.1	97	3	0	0	0	0.1	24	73	3	0	0	-0.2	61	36	3	0	0	0.0	0	58	39	3	0
Centre Plus Poll, 307564	-0.1	97	3	0	0	0	0.3	23	61	16	0	0	-0.2	58	39	3	0	0	-0.1	0	61	39	0	0
Centre Plus Poll, 9.183	0.1	85	15	0	0	0	0.2	30	56	14	0	0	0.4	18	56	26	0	0	0.1	0	44	56	0	0
Centre Plus WA Poll, 337919	0.1	83	17	0	0	0	0.1	31	66	3	0	0	0.1	38	52	7	3	0	-0.1	0	66	34	0	0
Hazeldean, 003542	0.0	90	7	3	0	0	-0.2	53	47	0	0	0	-0.3	63	37	0	0	0	-0.2	0	77	20	3	0
Kerin Poll, 130980	0.1	84	16	0	0	0	0.1	24	73	3	0	0	0.1	22	76	2	0	0	0.1	0	49	49	2	0
Mumblebone, 130389	-0.1	98	2	0	0	0	-0.5	80	20	0	0	0	0.2	28	62	10	0	0	0.2	3	35	60	2	0
Mumblebone, 130850	-0.1	100	0	0	0	0	0.1	32	63	5	0	0	0.2	24	71	5	0	0	0.3	0	29	71	0	0
Pastora Poll, 130011	0.1	87	11	0	0	2	-0.2	58	42	0	0	0	0.3	32	45	18	5	0	-0.2	2	74	24	0	0
Pastora Poll, 131634	0.0	93	7	0	0	0	-0.1	40	60	0	0	0	0.3	20	73	0	7	0	0.1	0	40	60	0	0
Pooginook Poll, 130083	-0.1	97	3	0	0	0	0.2	18	79	3	0	0	0.1	29	68	3	0	0	0.2	0	32	68	0	0
Rocklyn, 130022	-0.1	97	3	0	0	0	-0.2	55	45	0	0	0	-0.1	52	45	3	0	0	0.1	0	45	52	3	0
Roseville Park, 140019	0.1	77	23	0	0	0	0.1	29	68	3	0	0	-0.1	48	45	7	0	0	-0.1	0	68	32	0	0
Wattle Dale, 130115	-0.1	100	0	0	0	0	0.0	41	56	3	0	0	-0.1	48	44	8	0	0	-0.1	3	56	41	0	0
Willandra Poll, 120026	0.0	94	6	0	0	0	-0.2	53	44	3	0	0	-0.4	81	17	2	0	0	-0.3	6	72	19	3	0
Woodpark Poll, 130431	0.0	90	7	3	0	0	0.2	23	70	7	0	0	-0.2	53	43	4	0	0	-0.1	0	67	27	6	0
Avg.	1.1	92	8	0	0	0	1.7	38	58	4	0	0	1.7	42	50	6	2	0	2.5	1	55	43	1	0

^ **Historical Sire** evaluated under AMSEA's R&D project to validate the system of linkage in MERINOSELECT that has operated over the past 15-20 years. These sires were generally widely used 15-20 years ago and were selected for the R&D program based on their high ASBV accuracies. They demonstrate the progress the industry has made over that period.

Information on how to use the results in the table above can be found on page 18.

**Table 4b. Visual trait assessment – Wool quality and pigmentation**

Traits are reported as a deviation (Dev) from the average trait score for all progeny. The percentage of a sire's progeny assessed for each score is also reported. For the majority of breeder's objectives a negative deviation would be considered favourable and the larger the deviation the better.

Four pigmentation traits are reported as described on page 18. These are Fibre pigmentation, Non-fibre pigmentation, Recessive "black" and Random "spot". Fibre pigmentation and Non-fibre pigmentation are scored **1** to **5** however recessive black and random spot are scored **1** (no pigmentation of this type) or **5** (when the trait is expressed). Only the percentage scored 5 are reported for recessive black and random spot.

Breeder's flock, Sire number	Wool Quality												Pigmentation													
	Staple Weathering						Staple Structure						Fibre pigmentation						Non-fibre pigmentation						Black	Spot
	Dev	1	2	3	4	5	Dev	1	2	3	4	5	Dev	1	2	3	4	5	Dev	1	2	3	4	5	5	5
Billandri Poll, 121391	0.0	0	42	58	0	0	0.0	19	67	14	0	0	0.0	100	0	0	0	0	0.3	33	41	10	13	3	0	0
Bundilla, 120013	0.0	0	45	55	0	0	0.0	21	64	15	0	0	0.0	100	0	0	0	0	-0.3	60	31	9	0	0	0	0
Centre Plus Poll, 307564	-0.1	0	52	48	0	0	0.0	19	74	7	0	0	0.0	100	0	0	0	0	0.6	19	42	26	10	3	0	3
Centre Plus Poll, 9.183	0.1	4	33	56	7	0	0.1	26	44	30	0	0	0.0	100	0	0	0	0	0.0	26	65	9	0	0	0	0
Centre Plus WA Poll, 337919	-0.1	0	59	41	0	0	-0.1	28	59	13	0	0	0.0	100	0	0	0	0	-0.3	56	41	0	3	0	0	0
Hazeldean, 003542	-0.2	0	70	27	3	0	-0.1	23	73	4	0	0	0.0	100	0	0	0	0	-0.3	65	29	3	3	0	0	0
Kerin Poll, 130980	0.1	0	31	69	0	0	0.1	18	64	18	0	0	0.0	100	0	0	0	0	-0.4	60	36	4	0	0	0	0
Mumblebone, 130389	0.2	2	28	65	5	0	-0.2	32	62	6	0	0	0.0	100	0	0	0	0	0.1	43	26	29	2	0	0	0
Mumblebone, 130850	0.3	0	21	76	3	0	-0.5	55	45	0	0	0	0.0	100	0	0	0	0	0.0	32	57	8	3	0	0	0
Pastora Poll, 130011	-0.1	3	55	37	5	0	0.5	0	66	32	2	0	0.0	100	0	0	0	0	-0.3	57	35	8	0	0	0	0
Pastora Poll, 131634	0.2	0	33	53	14	0	0.2	7	73	20	0	0	0.0	100	0	0	0	0	0.2	38	31	25	6	0	0	6
Pooginook Poll, 130083	0.2	0	29	68	3	0	0.0	24	62	14	0	0	0.0	100	0	0	0	0	0.1	24	61	15	0	0	0	0
Rocklyn, 130022	0.0	0	48	48	4	0	-0.2	34	62	4	0	0	0.0	100	0	0	0	0	0.6	4	61	32	3	0	0	0
Roseville Park, 140019	-0.1	0	52	48	0	0	0.2	10	71	19	0	0	0.0	100	0	0	0	0	0.1	39	42	13	3	3	0	0
Wattle Dale, 130115	0.0	7	33	56	4	0	0.1	33	33	30	4	0	0.0	100	0	0	0	0	0.2	32	45	16	7	0	0	0
Willandra Poll, 120026	-0.3	17	47	31	2	3	0.0	31	50	19	0	0	0.0	100	0	0	0	0	-0.2	53	37	10	0	0	0	0
Woodpark Poll, 130431	-0.1	3	57	33	7	0	-0.1	27	63	10	0	0	0.0	100	0	0	0	0	-0.3	53	41	6	0	0	0	0
Avg.	2.6	3	43	51	3	0	1.9	24	61	15	0	0	1.0	100	0	0	0	0	1.8	41	42	13	3	1	0	0

^ **Historical Sire** evaluated under AMSEA's R&D project to validate the system of linkage in MERINOSELECT that has operated over the past 15-20 years. These sires were generally widely used 15-20 years ago and were selected for the R&D program based on their high ASBV accuracies. They demonstrate the progress the industry has made over that period. Information on how to use the results in the table above can be found on page 18.

**Table 4c. Visual trait assessments – Conformation**

Traits are reported as a deviation (Dev) from the average trait score for all progeny. The percentage of a sire's progeny assessed for each score is also reported. For the majority of breeder's objectives a negative deviation would be considered favorable and the larger the deviation the better. Face cover and body wrinkle are possible exceptions when for many breeders the optimum score is in the middle of the range.

Breeder's flock, Sire number	Conformation																													
	Jaw					Legs and Feet					Shoulder and Back					Face Cover					Body Wrinkle									
	Dev	1	2	3	4	5	Dev	1	2	3	4	5	Dev	1	2	3	4	5	Dev	1	2	3	4	5	Dev	1	2	3	4	5
Billandri Poll, 121391	0.0	100	0	0	0	0	-0.1	92	0	8	0	0	-0.1	97	0	3	0	0	0.1	0	0	86	14	0	0.1	17	64	11	8	0
Bundilla, 120013	0.0	100	0	0	0	0	-0.1	91	0	9	0	0	-0.1	100	0	0	0	0	-0.1	0	0	97	3	0	0.0	12	73	15	0	0
Centre Plus Poll, 307564	0.1	97	0	0	0	3	0.1	84	0	16	0	0	-0.1	97	0	3	0	0	-0.1	0	3	97	0	0	0.1	19	52	29	0	0
Centre Plus Poll, 9.183	0.0	100	0	0	0	0	-0.1	93	0	7	0	0	0.2	86	0	11	0	3	0.0	0	0	93	7	0	0.4	4	57	32	7	0
Centre Plus WA Poll, 337919	0.0	100	0	0	0	0	0.0	86	0	14	0	0	0.1	90	0	10	0	0	-0.1	0	3	90	7	0	0.2	10	66	17	7	0
Hazeldean, 003542	0.0	100	0	0	0	0	0.0	87	0	13	0	0	0.0	93	0	7	0	0	0.1	0	0	87	10	3	0.4	4	50	43	3	0
Kerin Poll, 130980	0.0	98	0	2	0	0	-0.2	96	0	4	0	0	0.1	89	0	11	0	0	-0.1	0	0	100	0	0	0.0	24	49	22	5	0
Mumblebone, 130389	0.0	98	0	2	0	0	0.3	72	0	28	0	0	-0.1	100	0	0	0	0	0.0	0	0	95	5	0	-0.5	55	42	3	0	0
Mumblebone, 130850	0.0	100	0	0	0	0	0.0	87	0	13	0	0	0.0	95	0	5	0	0	-0.1	0	3	92	5	0	-0.6	61	39	0	0	0
Pastora Poll, 130011	0.0	97	0	3	0	0	0.1	79	0	21	0	0	0.2	85	0	15	0	0	0.0	0	0	95	5	0	0.2	13	56	23	8	0
Pastora Poll, 131634	0.0	100	0	0	0	0	0.1	80	0	20	0	0	-0.1	100	0	0	0	0	-0.1	0	0	100	0	0	0.0	20	60	13	7	0
Pooginook Poll, 130083	0.0	97	0	3	0	0	-0.1	91	0	9	0	0	-0.1	97	0	3	0	0	0.1	0	0	82	18	0	-0.2	21	79	0	0	0
Rocklyn, 130022	0.0	100	0	0	0	0	-0.1	90	0	10	0	0	-0.1	97	0	3	0	0	0.2	0	0	79	17	4	-0.3	41	45	14	0	0
Roseville Park, 140019	0.0	97	0	3	0	0	0.0	87	0	13	0	0	0.0	94	0	6	0	0	-0.1	0	4	90	6	0	0.0	13	71	16	0	0
Wattle Dale, 130115	0.0	100	0	0	0	0	0.1	81	0	19	0	0	0.1	89	0	11	0	0	0.0	0	4	85	11	0	0.1	8	70	22	0	0
Willandra Poll, 120026	0.0	100	0	0	0	0	0.1	86	0	11	0	3	-0.1	100	0	0	0	0	0.2	0	0	75	22	3	0.0	25	53	17	5	0
Woodpark Poll, 130431	0.1	93	0	7	0	0	-0.2	97	0	3	0	0	0.1	90	0	10	0	0	0.0	0	0	90	10	0	0.0	20	60	13	7	0
Avg.	1.0	99	0	1	0	0	1.3	87	0	13	0	0	1.1	94	0	6	0	0	3.1	0	1	90	8	1	2.0	21	58	17	4	0

^ **Historical Sire** evaluated under AMSEA's R&D project to validate the system of linkage in MERINOSELECT that has operated over the past 15-20 years. These sires were generally widely used 15-20 years ago and were selected for the R&D program based on their high ASBV accuracies. They demonstrate the progress the industry has made over that period. Information on how to use the results in the table above can be found on page 18.

**Table 4d. Visual trait assessments – Breech**

Traits are reported as a deviation (Dev) from the average trait score for all progeny. The percentage of a sire's progeny assessed for each score is also reported. For the majority of breeder's objectives a negative deviation would be considered favorable and the larger the deviation the better. Breech scores were recorded at lamb marking.

Breeder's flock, Sire number	Breech																							
	Marking Breech Cover						Marking Breech Wrinkle						Marking Crutch Cover						Yearling Dag					
	Dev	1	2	3	4	5	Dev	1	2	3	4	5	Dev	1	2	3	4	5	Dev	1	2	3	4	5
Billandri Poll, 121391	-0.4	23	33	18	23	3	0.0	10	51	26	13	0	-0.4	23	33	18	23	3	Yearling Dag was not scored at classing, progeny had been crutched					
Bundilla, 120013	0.0	11	26	37	17	9	0.1	9	43	43	5	0	0.0	11	26	37	17	9						
Centre Plus Poll, 307564	-0.5	32	26	23	16	3	0.4	0	42	45	10	3	-0.5	32	26	23	16	3						
Centre Plus Poll, 9.183	-0.3	11	44	24	21	0	0.2	9	35	41	15	0	-0.3	11	44	24	21	0						
Centre Plus WA Poll, 337919	-0.2	12	41	19	25	3	0.3	3	38	53	3	3	-0.2	12	41	19	25	3						
Hazeldean, 003542	0.0	13	23	35	23	6	0.4	4	42	35	16	3	0.0	13	23	35	23	6						
Kerin Poll, 130980	0.3	4	22	31	36	7	0.1	13	40	33	11	3	0.3	4	22	31	36	7						
Mumblebone, 130389	0.1	14	17	33	29	7	-0.4	29	50	19	2	0	0.1	14	17	33	29	7						
Mumblebone, 130850	-0.1	18	30	15	32	5	-0.7	40	50	10	0	0	-0.1	18	30	15	32	5						
Pastora Poll, 130011	0.5	3	20	30	35	12	0.0	18	40	32	8	2	0.5	3	20	30	35	12						
Pastora Poll, 131634	0.0	25	25	19	6	25	-0.4	38	31	31	0	0	0.0	25	25	19	6	25						
Pooginook Poll, 130083	0.2	13	16	32	34	5	-0.4	23	53	24	0	0	0.2	13	16	32	34	5						
Rocklyn, 130022	-0.2	23	26	23	26	2	-0.2	13	61	23	3	0	-0.2	23	26	23	26	2						
Roseville Park, 140019	0.1	16	19	26	35	4	0.1	13	29	58	0	0	0.1	16	19	26	35	4						
Wattle Dale, 130115	0.0	6	26	45	19	4	0.5	4	32	42	16	6	0.0	6	26	45	19	4						
Willandra Poll, 120026	0.1	5	32	29	29	5	0.2	13	37	32	16	2	0.1	5	32	29	29	5						
Woodpark Poll, 130431	0.3	10	19	28	31	12	-0.2	12	59	25	4	0	0.3	10	19	28	31	12						
Avg.	2.8	14	26	27	26	7	2.4	15	43	34	7	1	2.8	14	26	27	26	7						

^ **Historical Sire** evaluated under AMSEA's R&D project to validate the system of linkage in MERINOSELECT that has operated over the past 15-20 years. These sires were generally widely used 15-20 years ago and were selected for the R&D program based on their high ASBV accuracies. They demonstrate the progress the industry has made over that period.

Information on how to use the results in the table above can be found on page 18.

Table 5. Sire means for measured traits

Sire means are the average performance of all the progeny of a sire. No account is made for factors that can improve the breeding value accuracy.

Ram code	Breeders flock, Ram number	No. of prog.	Ram averages for measured traits (deviations)											
			GFW %	CFW %	FD um	WT kg			Fat mm	EMD mm	FDCV %	Curv deg/mm	SL mm	SS N/ktex
			Y <sup>2</sup>	Y	Y	W	P	Y	Y	Y	Y	Y	Y	Y
1*	Billandri Poll, 121391	36	0.1	0.1	-0.2	0.7	0.1	-0.9	Not measured at time of publication	Not Measured at time of publication	0.3	4.2	-0.7	-4.9
2	Bundilla, 120013	33	0.0	0.0	0.3	0.0	0.3	0.1			-0.6	-0.3	2.1	3.5
3	Centre Plus Poll, 307564	31	0.3	0.2	-0.2	1.7	3.4	4.0			-0.6	-0.2	1.8	-0.6
4^	Centre Plus Poll, 9.183	27	0.1	0.0	0.4	-0.2	-0.8	0.0			0.0	3.4	-6.0	-0.7
5	Centre Plus WA Poll, 337919	29	0.1	0.0	-0.4	0.3	2.5	2.7			0.8	-0.3	1.4	0.3
6	Hazeldean, 003542	30	0.2	0.1	-0.7	-0.2	-1.8	-2.0			0.6	0.4	-3.3	-0.5
7	Kerin Poll, 130980	45	0.3	0.2	0.5	0.3	1.5	1.2			0.0	-4.5	6.5	3.6
8	Mumblebone, 130389	40	-0.4	-0.2	0.5	-2.3	-3.1	-3.4			-1.0	-3.1	4.7	-1.6
9	Mumblebone, 130850	38	-0.1	-0.1	0.3	0.3	1.8	1.7			-1.2	-2.2	6.7	2.1
10	Pastora Poll, 130011	38	-0.1	0.0	-0.3	0.1	-1.2	-2.2			0.5	2.9	-2.7	-0.1
11	Pastora Poll, 131634	15	-0.2	-0.1	-0.5	-0.1	0.7	0.6			0.6	3.3	-1.3	-2.6
12	Pooginook Poll, 130083	34	0.1	0.1	0.5	0.6	0.4	1.0			-0.1	-6.6	8.4	4.8
13	Rocklyn, 130022	29	-0.1	0.0	0.2	-0.4	-0.9	-1.5			-0.4	-2.9	2.3	2.5
14	Roseville Park, 140019	31	0.0	0.0	0.1	0.6	0.4	-0.2			0.0	-1.9	-1.8	0.6
15	Wattle Dale, 130115	27	-0.2	-0.1	-0.5	-0.3	-1.4	0.0			0.2	4.6	-5.8	-2.4
16*	Willandra Poll, 120026	35	-0.1	0.0	0.3	0.3	-1.0	-0.8			0.3	-0.1	-6.3	0.9
17	Woodpark Poll, 130431	29	-0.1	-0.2	-0.4	-1.4	-1.0	-0.3			0.5	3.4	-6.2	-5.0
Average performance		32	3.0 kg	2.0 kg	16.4 um	24.9 kg	33.2 kg	37.0 kg	mm	mm	18.8 %	90.5 deg/mm	78.9 mm	44.7 N/ktex

<sup>2</sup> W = Weaning (42 to 120 days); P = Post Weaning (120 to 300 days); Y = Yearling (300 to 400 days); H = Hogget (400 to 540 days); A = Adult (540 days and older).

<sup>^</sup> **Historical Sire** evaluated under AMSEA's R&D project to validate the system of linkage in MERINOSELECT that has operated over the past 15-20 years. These sires were generally widely used 15-20 years ago and were selected for the R&D program based on their high ASBV accuracies. They demonstrate the progress the industry has made over that period.

## Understanding the results

### Index Options – indexes reported on page 12.

Breeding Objective index options provide the relative value of sires based on a combination of the measured traits' genetic performance. The indexes used in this report are only some of the many indexes that can be used to describe an individual breeder's objective for measured traits.

**If a breeder is considering using a sire in this report it is critical to consider the performance of the breeder's flock relative to the performance standard in this report. The relative performance must be considered to establish the result that can be expected when a sire is used in a breeder's flock.**

All AMSEA site evaluation reports present 3 standard indexes to provide combined measured trait performance. These 3 AMSEA indexes are DP+; MP+; and FP+. These indexes are the same as MERINOSELECT indexes of that name however as there is no direct reproduction records captured by sire evaluation AMSEA do not include a Reproduction (NLW) FBV in their index calculations. As a result the 21% contribution by NLW in the DP+ index is not effectively applied by the index calculation.

### Index production system and breeding objectives

AMSEA  
DP+

**Dual Purpose:** Based on a meat focused production system where surplus progeny are sold as lambs and a portion of ewes are joined to terminal sires. Equal and high positive emphasis on CFW and meat traits (body weight, EMD). Maintain fibre diameter, staple strength and fat depth.




AMSEA  
MP+

**Merino Production:** Based on a balanced wool and meat production system where surplus progeny are sold as hoggets. Large increase in fleece weight. Moderate increase, staple strength, carcass traits and reproduction. Moderate reduction in fibre diameter.

AMSEA  
FP+

**Fibre Production:** Based on a wool focussed production system where wethers are retained, operating in an environment where worms cause economic losses. Large reduction in fibre diameter. Large increase in staple strength. Small reduction in WEC (if measured in the breeding program). Small increase in fleece weight. Little change in carcass traits and reproduction.

**Traits contribution to economic gain:** The percentage contribution of the traits listed to economic gain in a commercial flock that selects sires using the index.





### Accuracy of Flock Breeding Values

Flock Breeding Values (FBVs) are reported by Sheep Genetics (SG). FBVs express the expected performance of progeny of a sire relative to another sire in the evaluation when mated to the same standard of ewes. FBVs improve the accuracy of sire results because they account for the association between traits, adjustment for birth effects and the number of progeny a sire has in the analysis.

*True* Breeding Values would be achieved if the number of progeny evaluated for each sire were infinite. Because the number of progeny in the evaluation is not infinite, performance shown in this report is described as *Flock* Breeding Values.

Without progeny test information the correlation between the *Flock* and *True* Breeding Value of sires from different sources would be zero (0.0%). The correlation between *Flock* and *True* Breeding Value improves rapidly from 0.0% with no progeny to 77% with 10 progeny. The rate of improvement in correlation slows from 86% with 20 progeny, to 90% with 30 progeny and 92% with 40 progeny. With an infinite population the correlation is 100%. Note that the correlation used in the above example is for a trait such as fibre diameter with a high heritability (0.5).

A heritability of 0.5 indicates that half or 50% of the measured performance is passed onto offspring. A heritability of 0.35 indicates 35% is passed on. The FBVs that are shown in this report have already accounted for heritability and therefore describe the performance that can be expected from a sire's progeny.

### Link sires

Link sires provide the 'genetic link' between CTSE sites located across Australia to allow all sires entered in these site evaluations to have their performance reported relative to each other in *Merino Superior Sires*. *Merino Superior Sires* reports sires from across all effectively linked CTSE sites and across all evaluations at these sites. Link sires are therefore a vital component of the Central Test Sire Evaluation.

To be used as a link a sire must have at least 25 progeny assessed at 1st Assessment at one accredited site. Site reports provide valuable information not reported in *Merino Superior Sires* however *Merino Superior Sires* reports the performance of a large number of sires which can provide a wider perspective of the elite sires available across many flocks in Australia and New Zealand.

### Calculation – combined measured traits and combined visual trait performance

Combined measured trait performance is calculated as Index – 100. Three different index options are provided to cater for breeders' different breeding objectives.

Combined visual trait performance is calculated as:  
(Classer's Visual Grade Tops% - Culls%)/5, expressed as a deviation from the  
(average Tops% - average Culls%)/5

#### Example

Sires Performance: AMSEA DP+ Index Value = 119.7  
Tops % = 25.5 (average Tops% = 25.1)  
Culls% = 17.6 (average Culls% = 16.4)

Combined Measured = 119.7 – 100 = 19.7  
Combined Visual = ((25.5 - 17.6)/5) – ((25.1 – 16.4)/5)  
= 7.9/5 – 8.7/5 = 1.58 – 1.74 = -0.1

**Under the auspices of**  
**The Australian Merino Sire Evaluation Association**

